

## Movi Prep Preparation

### Central Texas Colon and Rectal Surgeons

David C. Fleeger, M.D. (512)452-9551 Fax (512)452-6997  
Ernest D. Graves III, M.D. (512)452-9551 Fax (512)452-6997  
Thiru Lakshman, M.D. (512) 452-9551 Fax (512) 452-6997  
Ricardo L. Solis M.D. (512)447-8911 Fax (512)447-8761  
John Mangione, M.D. (512)418-1979 Fax (512)418-1943  
William Robertson, M.D. (512)418-1755 Fax (512)418-1010  
Robert Cline, M.D. (512) 418-1763 Fax (512)372-9388

Name: \_\_\_\_\_

Your procedure has been scheduled for \_\_\_\_\_

at \_\_\_\_\_ Hospital.

Please be at admissions at \_\_\_\_\_.

This preparation includes a clear liquid diet and MoviPrep. Enemas and laxatives do not need to be taken or used unless prescribed. Do not eat seed foods or nuts for 48 hours prior to your procedure. Do not take Konsyl, Metamucil or other similar type fiber medications for 48 hours prior to your procedure.

Note: **Your prescriptions will be called in one week before your procedure.**

On \_\_\_\_\_ (the day prior to your procedure):

1. Begin clear liquid diet when you wake up.
2. At 12 PM place one pouch (labeled A) and one pouch (labeled B) in one of the two disposable containers, Add lukewarm drinking water to the top line of the container. Mix to dissolve. Place in refrigerator.

The MoviPrep container is divided by 4 marks.

2. At 4:00PM Every 15 minutes, drink the solution down to the next mark (approximately 8 oz.), until the full liter is consumed of the container.
3. Place the remaining pouch (labeled A) and the remaining pouch (labeled B) into the disposable container. Add lukewarm drinking water to the top line of the container. Mix to dissolve. Place in refrigerator
4. At 8:00PM Every 15 minutes, drink the solution down to the next mark (approximately 8 oz.), until the full liter is consumed of the container.

If you become nauseated or seem full, do not stop ingestion of the MoviPrep. Wait until the symptoms subside, then resume the process but drink 8 fluid ounces every 15 – 20 minutes (instead of every 10 minutes).

It is very important you finish the MoviPrep

5. **After midnight do not eat or drink anything** – you may use a minimum of water to take necessary medications.

#### Clear liquids permitted are listed below:

|   |  |
|---|--|
| Water   | Gatorade, Kool-Aid, lemonade, limeade  |
| Coffee – regular or decaffeinated (no milk or cream)      | Hawaiian or Delaware Punch             |
| Tea – iced, hot, or herbal                                | Broth – chicken or clear fat-free beef |
| Carbonated beverages (soft drinks) – regular or diet      | Bouillon or clear consommé             |
| Gelatin dessert – plain or fruit flavor (no red or green) | Snow cones or Popsicles                |
| Apple, grape, or cranberry juice                          | Hard candies, sugar and salt           |

No ice cream or orange juice with pulp.

## Special Instructions

Check with my medical assistant concerning the medicines you take daily. **Stop all aspirin, aspirin-containing medications, nonsteroidal anti-inflammatory drugs, and Ibuprofen medications seven (7) days prior to your colonoscopy.** If you are on Coumadin (Warfarin), Plavix or other blood thinners, you will be given special instructions by the doctor. Tylenol and Tylenol-containing pain medications are fine.

**All patients must have an adult available to accompany or drive them home.** Patients cannot use a public conveyance (taxi or bus) without an adult to accompany them. We ask that you have a family member or friend present at the conclusion of the procedure, so the doctor can discuss the results with them. You will stay in the Recovery Area after your colonoscopy is completed, until you are safely capable of leaving. Bring a book!

**OUR OFFICE HOURS ARE M-F FROM 9-5PM. IF AN EMERGENCY DOES ARISE AND OUR OFFICE IS CLOSED PLEASE CALL THE OFFICE AND THE RECORDING WILL INSTRUCT YOU ON HOW TO REACH THE DOCTOR.**

**Contact your physician if you experience abdominal pain, fever, rectal bleeding, persistent vomiting, or other problems during or following the colonoscopy preparation.**

If you have had blood drawn, X-rays or EKG performed recently, please advise the office, as this may prevent redundant lab tests.

No jewelry should be worn, wear comfortable clothes, and bring your insurance cards.

If you have any difficulty with your blood clotting, please notify the office (i.e. do you bruise easily? do your cuts bleed longer than other people's cuts? etc.)

## Home Care Instructions After Your Colonoscopy

**Activity Level: All patients must have an adult to accompany them home.** You may feel sleepy for several hours. Your planned activities following your hospital departure should be limited accordingly. Absolutely do not drive a car or operate machinery on the day of your colonoscopy. Resume normal activities on the day following your colonoscopy.

**Diet:** After your colonoscopy is over you may eat that day, but do not eat any extra spicy foods, or foods that are hard to digest. A regular diet may be eaten the following day. If polyps were removed (or biopsies taken). Do not take aspirin, aspirin-containing medications or Ibuprofen for at least two (2) weeks. Do not resume blood thinners until you have discussed it with your doctor.

**Bowel Movements:** Your normal regularity may not return for 2-3 days following your colonoscopy. Small amounts of bright red blood, especially on toilet tissue, is not an unusual occurrence.

**Report to your physician any episodes of:**

1. Bright red bloody bowel movements;
2. Any persistent severe abdominal pain;
3. Temperature elevation greater than 100.5 degrees;
4. Persistent nausea and/or vomiting.

**Return appointments and biopsy reports:** If polyps are removed during your colonoscopy, we will call within the next 48 – 72 hours with your final report status after the pathologist has reported their findings. If you have not heard from us in a week, please call the office. However, it is very rare that the initial impression given to you at the time of your procedure differs from the final report. We will also inform you of any other laboratory or X-ray results, which are available at that time, and we will discuss the need for any additional or follow-up appointments.